



UNIVERSITI PUTRA MALAYSIA

**URINARY PURINE DERIVATIVES EXCRETION AS A METHOD FOR
ESTIMATION OF RUMEN MICROBIAL PROTEIN PRODUCTION IN
SWAMP BUFFALOES AND ZEBU CATTLE**

OPART PIMPA

FP 2002 7

**URINARY PURINE DERIVATIVES EXCRETION AS A METHOD FOR
ESTIMATION OF RUMEN MICROBIAL PROTEIN PRODUCTION IN
SWAMP BUFFALOES AND ZEBU CATTLE**

OPART PIMPA

**DOCTOR OF PHILOSOPHY
UNIVERSITI PUTRA MALAYSIA**

2002



**URINARY PURINE DERIVATIVES EXCRETION AS A METHOD FOR
ESTIMATION OF RUMEN MICROBIAL PROTEIN PRODUCTION IN
SWAMP BUFFALOES AND ZEBU CATTLE**

By

OPART PIMPA

**Thesis Submitted to the Graduate School, Universiti Putra Malaysia, in Fulfilment
of the Requirements for the Degree of Doctor of Philosophy**

January 2002



I wish to dedicate this thesis to my beloved and respected parents, teachers and supervisors who had enriched my knowledge and experience. I also wish to dedicate this thesis to all the buffaloes and cattle used for my experiments.

Abstract of the thesis presented to the Senate of Universiti Putra Malaysia in
fulfilment of the requirement for the degree of Doctor of Philosophy

**URINARY PURINE DERIVATIVES EXCRETION AS A METHOD FOR
ESTIMATION OF RUMEN MICROBIAL PROTEIN PRODUCTION IN
SWAMP BUFFALOES AND ZEBU CATTLE**

By

OPART PIMPA

January 2002

Chairman: Associate Professor Liang Juan Boo, Ph.D.

Faculty: Agriculture

Prediction equations based on urinary purine derivatives (PD) excretion rate as an index to predict rumen microbial protein production have been developed for European cattle and sheep. However, there is evidence to suggest that those equations may not be applicable directly to tropical swamp buffaloes (*Bubalus bubalis*) and zebu cattle (*Bos indicus*). To establish similar equations for the above two species of ruminant, five studies were conducted. In the first study, endogenous PD excretion rate determined by fasting procedure for swamp buffaloes and the Malaysian indigenous KK cattle (zebu cattle) were 199 and 300 $\mu\text{mol/kg}^{0.75}$ /day, respectively. Urinary PD excretion rate per kg digestible organic matter intake (DOMI) for buffaloes (8.19 mmol/kg DOMI) was significantly lower than that for KK cattle (15.45 mmol/kg DOMI). The second study examined the relationship between daily urinary PD excretion (Y , mmol) and exogenous purine bases (PB) supply via duodenal infusion (X , mmol/day). The relationship obtained were $Y = 0.12X + 12.78$ ($r^2 = 0.45$) for buffaloes and $Y = 0.85X + 7.15$ ($r^2 = 0.62$) for KK cattle, suggesting that 12% and 85% of the supplied exogenous purine were excreted in the urine of buffaloes and

zebu cattle, respectively. In the third study, labelled $[8-^{14}\text{C}]$ uric acid marker was used to test the hypothesis that the lower recovery rate of urinary PD in swamp buffaloes was due to their higher recycling of plasma PD as compared to KK cattle. The averaged non-renal PD loss of plasma PD for swamp buffaloes and KK cattle did not differ significantly. The result thus rejected the above hypothesis. A follow-up study was conducted to examine whether the lower urinary PD excretion rate for buffaloes could be due to lower absorption in the small intestine as compared to cattle. The results indicated that purine absorption rate for buffaloes as percent of disappearance of the first segment of small intestine was not significantly different from that of zebu cattle. However a significantly larger quantity of digesta and PB was noted in the last 3 segments of the small intestine of buffaloes than of cattle. The above results suggested that the lower urinary PD excretion rate in the buffaloes could be partially due to their lower absorption rate in the small intestine as compared to cattle. Based on the endogenous and PD excretion rate obtained in this thesis, the relationships between daily urinary PD excretion (Y , mmol) and daily microbial purine supply (X , mmol) were $Y = 0.12X + 0.20 \text{ kg}^{0.75}$ for buffaloes and $Y = 0.85X + 0.30 \text{ kg}^{0.75}$ for zebu cattle. The equation for zebu cattle is similar to those published for European cattle. Due to the extremely low recovery rate (12%), the proposed equation for buffaloes should be used with great caution. Based on the above equations, microbial N yield for buffaloes and zebu cattle can be calculated from the following equation;

$$\text{Microbial N (gN/day)} = \frac{X (\text{mmol/day}) \times 70}{0.073 \times 0.83 \times 1000} = 1.16 X$$

In the above equation, digestibility of microbial purine was assumed to be 0.83 and N content of purine was 70 mg N/mmol. The ratio of purine N: total N in mixed rumen microbes was taken as 7.3: 100 for buffaloes and zebu cattle.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk Ijazah Doktor Falsafah

**TERBITAN PURINA URINA SEBAGAI SATU KAEDAH UNTUK
MENGANGGARKAN PENGHASILAN PROTEIN MIKROB RUMEN KERBAU
DAN LEMBU ZEBU**

Oleh

OPART PIMPA

Januari 2002

Pengerusi: Prof. Madya Dr. Liang Juan Boo

Fakulti : Pertanian

Kadar pengeluaran terbitan purina dalam urina (PD) telah digunakan sebagai indeks bagi menganggar penghasilan protein mikrob rumen ternakan ruminan. Persamaan dihasilkan untuk lembu dan bebiri Eropah tetapi ia tidak boleh digunakan secara langsung pada kerbau (*Bubalus bubalis*) dan lembu (*Bos indicus*) tropikal. Oleh itu, lima eksperimen telah dijalankan bagi mendapatkan persamaan untuk kerbau dan lembu. Dalam eksperimen pertama, kadar pengeluaran PD endogenus untuk kerbau dan lembu KK yang ditentukan dengan prosedur berpuasa adalah $199 \mu\text{mol/kg}^{0.75}/\text{hari}$, dan $300 \mu\text{mol/kg}^{0.75}/\text{hari}$, masing-masing. Kadar pengeluaran PD urin per kg bahan organik hadaman (DOMI) kerbau ($8.19 \text{ mmol/kg DOMI}$) didapati lebih rendah ($P < 0.05$) dari lembu KK ($15.45 \text{ mmol/kg DOMI}$). Eksperimen yang kedua mengkaji pertalian di antara pengeluaran PD urin harian (Y , mmol) dan bes purina eksogenus (PB) yang di suntik melalui infusi duodenal (X , mmol/hari). Pertalian yang didapati bagi kerbau adalah $Y = 0.12X + 12.78$ ($r^2 = 0.45$) manakala bagi lembu KK adalah $Y = 0.85X + 7.15$ ($r^2 = 0.62$), mencadangkan bahawa 12% purina eksogenus yang dibekalkan dikeluarkan oleh kerbau,

sementara lembu mengeluarkan 85%. Dalam kajian ketiga, asid urik berlabel ($8-^{14}\text{C}$) digunakan bagi menguji hipotesis yang menyatakan bahawa kadar pemulihan bagi PD urina kerbau disebabkan kitaran semula plasma PD yang lebih tinggi berbanding lembu KK. Purata nilai kehilangan bukan-renal untuk kerbau dan lembu tidak berbeza. Oleh itu, keputusan ini menolak hipotesis tersebut. Kajian seterusnya dijalankan untuk meneliti samada kadar pengeluaran PD yang rendah pada kerbau disebabkan oleh penyerapan purina yang rendah di usus kecil kerbau. Hasil kajian menunjukkan bahawa kadar penyerapan purina kerbau tidak berbeza dengan lembu. Walaupun begitu, kuantiti PD dalam digesta kerbau didapati lebih tinggi berbanding lembu pada tiga bahagian akhir usus kecil. Keputusan ini mencadangkan bahawa perbezaan di antara kedua spesies ruminan ini mungkin disebabkan oleh kadar penyerapan purina yang rendah di usus kecil kerbau. Berdasarkan keputusan tersebut, pertalian di antara pengeluaran harian PD dalam urin (Y , mmol) dan bekalan harian purina mikrobial (X , mmol) bagi kerbau adalah $Y = 0.12X + 0.20 \text{ kg}^{0.75}$ dan bagi lembu adalah, $Y = 0.85X + 0.30 \text{ kg}^{0.75}$. Berdasarkan kedua persamaan di atas, penghasilan N mikrob boleh dianggarkan dengan persamaan di bawah:

$$\text{Mikrobial N (gN/hari)} = \frac{X (\text{mmol/hari}) \times 70}{0.073 \times 0.83 \times 1000} = 1.16 X \text{ bagi kerbau dan lembu}$$

Persamaan ini menganggap penghadaman purina mikrobial adalah 0.83 dan kandungan N purina adalah 70 mg N/mmol. Nisbah kadar N purina : N keseluruhan dalam campuran mikrob rumen diambil sebagai 7.3 : 100 bagi kerbau dan lembu. Persamaan bagi pembekalan purin mikrobial dengan menggunakan pengeluaran PD dalam urin untuk lembu KK adalah sama dengan persamaan untuk lembu Europah, Oleh kerana kadar kedapatan semula untuk kerbau (12%) sangat rendah maka, persamaan ini perlu digunakan secara berhati-hati.

ACKNOWLEDGEMENTS

I would like to express my deepest appreciation and sincere gratitude to my supervisor, Associate Professor Dr. Liang Juan Boo for his invaluable guidance, constructive criticisms, and constant encouragement throughout the course of my study. My appreciation and gratitude also go to Associate Professor Dr. Zainal Aznam Bin Mohd. Jelani and Associate Professor Dr. Norhani Bt. Abdullah, members of my Supervisory Committee for their invaluable guidance and suggestions.

I wish to express my deepest appreciation to Professor Dr. Metha Wanapat for encouraging me to pursue this study. I would like to thank my fellow students, particularly Mr. Thongsuk Jetana, Mr. Bodee Khamseekhiew and Miss Latiffah Zakaria for their help in accomplishing this thesis. I would like to thank Dr. M.Y. Loqman, Dr. G.K. Mohd Azam Khan and many of the veterinary students for their help in preparing the rumen-duodenal fistulated animals for my studies. Thanks are also due to the technicians of the Department of Animal Science. I also like to thank Dr. Nashriyah of the Malaysian Institute for Nuclear and Technology (MINT) for her assistance in the tracer analysis, and Dr. Jothi Malar Panamdham and Mr. Ahmad Shokri b. Othman for their assistance in statistical analysis.

I specially thank the Director, Livestock Research Center of MARDI and the Head of Department of Animal Science, UPM for providing the research facilities. I am grateful to the Ministry of Science, Technology and Environment, Malaysia and the Joint FAO/IAEA Division of Nuclear Techniques in Food and Agriculture, International

Atomic Energy Agency, Vienna, for jointly financing my study. Appreciation goes to Naresuan University, Ministry of University Affairs, Kingdom of Thailand for granting me the study leave.

My special thanks to my beloved wife Benchamaporn Pimpa for her inspiration, moral encouragement and sacrifices during the course of this study. Finally, I am grateful to the cattle and buffaloes used in this study.

I certify that an Examination Committee met on 15th January 2002 to conduct the final examination of Opart Pimpa on his Doctor of Philosophy thesis entitled "Urinary Purine Derivatives Excretion as a Method for Estimation of Rumen Microbial Protein Production in Swamp Buffaloes and Zebu Cattle" in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The Committee recommends that the candidate be awarded the relevant degree. Members of Examination Committee are as follows:

HO YIN WAN, Ph.D.

Professor
Faculty of Science and Environmental Studies
Universiti Putra Malaysia
(Chairperson)

LIANG JUAN BOO, Ph.D.

Associate Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Member)

ZAINAL AZNAM BIN MOHD. JELAN, Ph.D.


Associate Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Member)

NORHANI ABDULLAH, Ph.D.

Associate Professor
Faculty of Science and Environmental Studies
Universiti Putra Malaysia
(Member)

JOAQUIM BALCELLS, Ph.D.

Faculty de Veterinaria
Universidad de Zaragoza
Spain
(Independent Examiner)




AINI IDERIS, Ph.D.

Professor,
Dean of Graduate School,
Universiti Putra Malaysia

Date: 14 FEB 2002

The thesis submitted to the Senate of Universiti Putra Malaysia has been accepted as fulfilment of the requirement for the degree of Doctor of Philosophy.




AINI IDERIS, Ph.D.
Professor/Dean of Graduate School
Universiti Putra Malaysia

Date:

DECLARATION

I hereby declare that the thesis is based on my original work except for quotations and citations which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at UPM or other institutions.

A handwritten signature in black ink, appearing to read 'part' followed by 'impa' with a large loop at the end.

OPART PIMPA

Date : 14 February 2002

TABLE OF CONTENTS

	Page
ABSTRACT.....	iii
ABSTRAK.....	v
ACKNOWLEDGEMENTS.....	vii
APPROVAL SHEETS.....	ix
DECLARATION FORM.....	xi
LIST OF TABLES.....	xviii
LIST OF FIGURES.....	xxi
LIST OF ABBREVIATIONS.....	xxiii
CHAPTER	
1 INTRODUCTION.....	1
2 LITERATURE REVIEW.....	4
2.1 Buffaloes and Cattle Production in Malaysia.....	4
2.2 Comparative Efficiency of Feed Utilization by Buffaloes and Cattle.....	5
2.3 Feed Resources.....	7
2.4 Quantification of Ruminal Microbial Protein Synthesis.....	8
2.4.1 Internal Microbial Markers.....	8
2.4.2 External Microbial Markers.....	10
2.5 Estimation of Digesta Flow.....	11
2.6 Nucleic Acid (NA) Digestion in the Rumen.....	12
2.7 The Ratio of Rumen Microbial Nitrogen and Purine Nitrogen...	13
2.8 Digestion and Absorption of Microbial Nucleic Acids in the Small Intestine.....	15
2.8.1 The Nucleic Acid Metabolite Products in the Blood and Urine.....	17
2.9 Use of ¹⁴ C Isotopes for the utilization of Microbial Nucleic Acid	18
2.9.1 Use Isotopes (¹⁴ C and ¹⁵ N) to Study Recovery Rate of Purine Derivatives and Glomerular Filtration Rate (GFR).....	19
2.10 Purine Anabolism.....	20
2.10.1 Purine Synthesis <i>de novo</i>	20
2.10.2 Purine Salvage.....	21
2.11 The Xanthine Oxidase Activity in Ruminants.....	22
2.12 The Use of Purine Derivatives in Measuring Microbial Protein Synthesis.....	23
2.13 Purine Derivatives of Endogenous Origin.....	26
2.14 Difference between the Endogenous and Exogenous PD.....	29



2.15	Urinary PD Components in Buffaloes and Cattle.....	31
2.16	The Relationship of Urinary Purine Derivatives and Feed Intake.....	33
2.17	Recovery Rate of Exogenous Nucleic Acid (NA) Supply to Urinary Purine Derivatives.....	36
2.18	Prediction of Rumen Microbial Outflow Based on Urinary PD Excretions.....	39
2.19	Plasma PD in Relation to Urinary PD.....	41
2.20	Creatinine Excretion in Ruminants.....	43
2.21	Glomerular Filtration Rate (GFR).....	46
2.22	Use of the Purine Derivatives : Creatinine (PD: C) in Spot Urine Samples for Estimating of Rumen Microbial Yield.....	50
2.23	The Saliva and Milk Purine Derivatives.....	51
3	ESTIMATION OF BASAL PURINE EXCRETION AND RESPONSE OF PURINE EXCRETION TO FEED INTAKE	54
3.1	Introduction.....	54
3.2	Experiment 1: Urinary Excretion of Endogenous Purine Derivatives (PD).....	55
3.2.1	Materials and Methods.....	55
3.2.1.1	Animals and Their Management.....	55
3.2.1.2	Sample Collection.....	56
3.2.1.3	Chemical Analysis.....	57
3.2.1.3.1	Urine and Blood plasma Preparation for PD Analysis	57
3.2.1.3.2	HPLC Instrumentation and Chromatographic Conditions.....	57
3.2.1.4	Calculation of Glomerular Filtration Rate (GFR).....	60
3.2.1.5	Statistical Analysis.....	60
3.2.2	Results.....	60
3.2.2.1	Animal Condition.....	60
3.2.2.2	Endogenous Purine Derivatives (PD) Excretion.	61
3.2.2.3	Glomerular Filtration Rate (GFR)	63
3.2.3	Discussion.....	63
3.2.3.1	Endogenous Purine Derivatives (PD)	63
3.2.3.2	Creatinine and Glomerular Filtration Rate (GFR).....	66
3.3	Experiment 2: Urinary Excretion of Purine Derivatives in Response to Different Levels of Feed Intake.....	67
3.3.1	Materials and Methods.....	67
3.3.1.1	Animals and Feeding.....	67
3.3.1.2	Sample Collection and Preparation.....	68
3.3.1.3	Chemical Analysis.....	71

3.3.1.4	Calculation of Glomerular Filtration Rate (GFR), Tubular Load (TL) and Reabsorption (RB).....	72
3.3.1.5	Statistical Analysis.....	74
3.3.2	Results.....	74
3.3.2.1	Digestibility.....	74
3.3.2.2	Urinary PD Excretion Rate.....	75
3.3.2.3	The Creatinine Excretion.....	75
3.3.2.4	Urinary Nitrogen Excretion.....	77
3.3.2.5	The Relationships between Urinary PD Excretion Rates and DOMI.....	77
3.3.2.6	Plasma Purine Derivatives.....	79
3.3.2.7	The Glomerular Filtration Rate (GFR), Tubular Load (TL) and Reabsorption (RB).....	79
3.3.3	Discussion.....	82
3.3.3.1	Dry Matter Intake (DMI) and Digestibility.....	82
3.3.3.2	The Response of Purine Derivatives Excretion to Feed Intake.....	82
3.3.3.3	Creatinine Excretion.....	83
3.3.3.4	Urinary Nitrogen Excretion.....	83
3.3.3.5	Glomerular Filtration Rate (GFR).....	84
3.3.4	Conclusion.....	84
4	RECOVERY RATE OF PURINE DERIVATIVES IN URINE OF KEDAH- KELANTAN CATTLE AND SWAMP BUFFALOES AFTER DUODENALLY INFUSED WITH INCREMENTAL AMOUNT OF PURINE BASES.....	85
4.1	Introduction.....	85
4.2	Experiment 1: Urinary Excretion Rate of Duodenal Purine Derivatives in Kedah- Kelantan Cattle.....	86
4.2.1	Materials and Methods.....	86
4.2.1.1	Animals and Their Management.....	86
4.2.1.2	Experimental Design.....	88
4.2.1.3	Measurement of Basal Purine in Duodenal Digesta.....	90
4.2.1.4	Purine Bases (PB) Infusion.....	90
4.2.1.5	Urine and Blood Sampling.....	91
4.2.1.6	Chemical Analysis.....	93
4.2.1.6.1	Purine derivatives Analysis.....	93
4.2.1.6.2	Co and Yb Marker Analysis.....	93
4.2.1.6.3	Purine bases Analysis.....	94
4.2.1.7	Duodenal PB Flow Rate Calculation.....	95
4.2.1.8	Calculation of Glomerular Filtration Rate (GFR), Tubular Load (TL) and Reabsorption (RB).....	97
4.2.1.9	Data and Statistical Analysis.....	97
4.2.2	Results.....	98

4.2.2.1	Condition of Cattle.....	98
4.2.2.2	Basal Purine in Duodenal Digesta.....	98
4.2.2.3	Purine Derivatives Recovery.....	101
4.2.2.4	Purine Derivatives Excretion in Relation to Purine Supply.....	101
4.2.2.5	The Concentration of PD in Plasma.....	101
4.2.2.6	Glomerular Filtration Rate (GFR), Tubular Load (TL) and Reabsorption (RB).....	103
4.2.3	Discussion.....	103
4.2.3.1	Basal Purine in Duodenal Digesta.....	103
4.2.3.2	Purine Derivatives Recovery.....	105
4.2.3.3	The Relationship between Total PD Excretion and PB Absorption	105
4.2.3.4	The Concentration of PD in Plasma.....	107
4.2.3.5	Glomerular Filtration Rate (GFR), Tubular Load (TL) and Reabsorption (RB).....	107
4.3	Experiment 2: Urinary Purine Derivatives in Swamp Buffaloes after Purine Bases Infusion into the Duodenum.....	108
4.3.1	Materials and Methods.....	108
4.3.1.1	Purine Bases (PB) Infusion.....	110
4.3.1.2	Urine and Blood Sampling.....	112
4.3.1.3	Analytical Methods.....	112
4.3.1.4	Calculation of Glomerular Filtration Rate (GFR), Tubular Load (TL) and Reabsorption (RB).....	113
4.3.1.5	Satatistical Analysis.....	113
4.3.2	Results.....	113
4.3.2.1	Condition of Buffaloes.....	113
4.3.2.2	Purine Derivatives Recovery.....	114
4.3.2.3	The Concentration of Plasma PD and Creatinine.....	114
4.3.2.4	Glomerular Filtration Rate (GFR), Tubular Load (TL) and Reabsorption (RB).....	118
4.3.3	Discussion.....	118
4.3.3.1	The Urinary PD and Creatinine Excretion.....	118
4.3.3.2	The Relationship between Total Urinary PD Excretion and PB Infusion.....	121
4.3.3.3	The Concentration of PD and Creatinine in Plasma.....	122
4.3.3.4	Glomerular Filtration Rate (GFR), Tubular Load (TL) and Reabsorption (RB).....	123
4.3.4	Conclusions.....	125
5	RECOVERY RATE OF PLASMA PURINE DERIVATIVES IN THE URINE OF SWAMP BUFFALOES AND KEDAH- KELANTAN CATTLE BY TRACER STUDIES.....	126
5.1	Introduction.....	126

5.2	Experiment 1: Recovery of Plasma Purine Derivatives in the Urine of Swamp Buffaloes and Kedah-Kelantan Cattle at Two Levels of Feed Intake.....	127
5.2.1	Materials and Methods.....	127
5.2.1.1	Sample Collection and Preparation.....	128
5.2.1.2	Analytical Methods.....	129
5.2.1.3	Data Analysis.....	129
5.2.2	Results.....	130
5.2.2.1	Plasma Isotope Dilution	130
5.2.2.2	Recovery of Tracer in Urine.....	134
5.2.3	Discussion.....	138
5.2.3.1	Plasma ¹⁴ C PD Degradation.....	138
5.2.3.2	The Recovery of Labeled ¹⁴ C PD in Urine.....	138
5.3	Experiment 2: Recovery of Plasma Purine Derivatives in Swamp Buffaloes with Duodenal Purine Bases Infusion.....	140
5.3.1	Materials and Methods.....	140
5.3.1.1	Blood Plasma and Urine Sampling.....	141
5.3.1.2	Analytical Methods.....	141
5.3.1.3	Data Analysis.....	142
5.3.2	Results	142
5.3.2.1	Plasma Isotope Dilution	142
5.3.2.2	Isotope Dilution in Urine.....	142
5.3.3	Discussion.....	146
5.3.3.1	The Plasma Isotope Dilution Curves.....	146
5.3.3.2	Recovery rate of ¹⁴ C PD in Urine.....	149
5.3.4	Conclusion.....	150
6	ABSORPTION OF NUCLEIC ACID IN THE SMALL INTESTINES OF SWAMP BUFFALOES AND KEDAH-KELANTAN (KK) CATTLE	152
6.1	Introduction.....	152
6.2	Materials and Methods.....	153
6.2.1	Animals and Feeding.....	153
6.2.2	Collection of Digesta.....	154
6.2.3	Analytical Methods.....	154
6.2.3.1	Determination of DM and Cr ₂ O ₃ Marker in Digesta.....	154
6.2.3.2	RNA and DNA Analysis.....	156
6.2.3.3	Purine Bases Analysis.....	157
6.2.3.4	Data Analysis.....	157
6.3	Results.....	158
6.3.1	Concentration of RNA and DNA.....	158
6.3.2	Concentration of Purine Bases in Digesta.....	160
6.4	Discussion.....	168
6.5	Conclusions.....	171

7	PURINE NITROGEN: TOTAL NITROGEN CONTENTS OF MIXED RUMEN MICROBIAL POPULATION IN SWAMP BUFFALOES AND KEDAH-KELANTAN CATTLE.....	172
7.1	Introduction.....	172
7.2	Materials and Methods.....	173
7.2.1	Animal and Their Management.....	173
7.2.2	Experimental Methods.....	173
7.2.3	Sample Preparation.....	174
7.2.4	Samples Analysis.....	176
7.2.5	Data Analysis.....	176
7.3	Results.....	176
7.3.1	Chemical Composition of Rumen Microbial Extracts.....	176
7.4	Discussion.....	178
7.5	Conclusion.....	180
8	GENERAL DISCUSSION AND CONCLUSION.....	182
	BIBLIOGRAPHY.....	187
	VITA.....	204

LIST OF TABLES

Table		Page
2.1	Purine-N: total-N ratios (%) in the rumen fluid-associated bacteria (FB) and rumen fluid-associated microbes (FM), the particle-associated bacteria (PAB) and particle-associated microbes (PAM) in the rumen and digesta liquid-associated bacteria (LAB) of ruminants fed on various diets of various roughage: concentrate (Con).....	14
2.2	Xanthine oxidase activities in plasma $\mu\text{mol}/\text{min}/\text{L}$ (unit/L), fresh liver tissue (unit/g) and intestinal (unit/g wet mucosa cell)	24
2.3	Endogenous purine derivatives in different species ($\mu\text{mol}/\text{kg}^{0.75}/\text{day}$).....	28
2.4	The proportion (%) of purine derivatives in different species and experiments.....	32
2.5	The relationship between urinary purine derivatives excretion (mmol/day) and digestible organic matter intake (kg, DOMI) in difference species and experiments.....	35
2.6	The recovery of urinary purine derivatives by duodenal infusion of purines.....	38
2.7	The prediction equation for rumen microbial outflow based on urinary purine derivatives excretion in different species.....	40
2.8	The creatinine excretion ($\mu\text{mol}/\text{kg}^{0.75}/\text{day}$) in different species..	45
2.9	The glomerular filtration rate (GFR, L/day) in ruminants	49
3.1	Daily urinary excretion of purine derivatives (PD), GFR and PD tubular load in swamp buffaloes and cattle during pre-fasting period.....	64
3.2	Daily urinary excretion of purine derivatives (PD), creatinine, GFR of PD in swamp buffaloes and cattle during fasting period.....	65
3.3	Chemical composition of the ingredients used in feeding trial...	69
3.4	Treatment levels for feeding trial (kg DM/day).....	70

3.5	Effect of feed intake levels on dry matter (DM) and organic matter (OM) digestibility (%), digestible DM intake (kg/day) and digestible OM intake (kg/day) in swamp buffaloes and KK cattle.....	73
3.6	Daily PD excreted, creatinine ($\text{mmol/kg}^{0.75}$) and urine nitrogen ($\text{mg/kg}^{0.75}$) of swamp buffaloes and KK cattle in different feeding levels.....	76
3.7	Mean blood plasma PD and creatinine (mmol/L) of swamp buffaloes and KK cattle with different feeding levels...	80
3.8	Effect of feed intake on glomerular filtration rate (GFR, L/day) tubular load (TL, mmol/day) and net reabsorb (RB, mmol/day) of PD in swamp buffaloes and KK cattle	81
4.1	Estimated daily flow rate of digesta, purine bases (PB) concentration in digesta and daily PB flow*.....	99
4.2	Daily excretion of urinary PD, creatinine, urine nitrogen and recovery rates as derivatives in urine in the different treatments in KK cattle.....	102
4.3	PD and creatinine concentrations in plasma, glomerular filtration rate (GFR), tubular load (TL) and net re-absorption (RB) of purine in KK cattle in different treatments.....	104
4.4	Daily excretion of urinary PD and creatinine (mmol/day) in swamp buffaloes after duodenally infusion with five levels of purine bases (mmol/day).....	115
4.5	Daily excretion of urinary PD and creatinine ($\text{mmol/kg}^{0.75}$) in swamp buffaloes after duodenally infusion with five levels of purine bases (mmol/day).....	117
4.6	Plasma PD (mmol/L) in swamp buffaloes in different treatments of exogenous purine bases infusion (mmol/day)...	119
4.7	Glomerular filtration rate (GFR, L/day), tubular load (TL) and reabsorption (mmol/day) in swamp buffaloes after duodenally infusion with five levels of purine bases, PB (mmol/day).....	120
5.1	The plasma ^{14}C tracer PD (allantoin plus uric acid) recovery in different time after tracer injection (μCi and in % of total injection) in swamp buffaloes and KK cattle.....	133

5.2	Urinary tracer $^{14}\text{C}_1$ (allantoin plus uric acid and other form) recovery ($\% \pm \text{SE}$) in swamp buffaloes and KK cattle.....	135
5.3	Total urinary ^{14}C tracer recovery (%) in swamp buffaloes and KK cattle.....	137
5.4	Plasma ^{14}C tracer PD (allantoin plus uric acid) concentration (dpm/mL) in swamp buffaloes	143
5.5	Total urinary ^{14}C tracer PD (allantoin plus uric acid) recovery (%) in swamp buffaloes at different time after tracer ^{14}C uric acid injection.....	145
5.6	The urinary ^{14}C tracer recovery ($\% \pm \text{SEM}$) in swamp buffaloes.....	148
6.1	RNA and DNA concentrations (mg/mg Cr_2O_3) and absorption (% of disappearance from the first segment) in digesta from swamp buffalo and KK cattle at different segments of the small intestines (values are measured of 3 animals with their respective standard errors).....	159
6.2	Concentrations ($\mu\text{mol/mg}$ Cr_2O_3) of purine bases in digesta from different segments of swamp buffalo and KK cattle small intestines (values are measured of 3 animals with their respective standard errors).....	164
6.3	Percentage absorption (% disappearance from the first segment) of purine bases in digesta from different intestine segments of swamp buffalo and KK cattle (values are measured of 3 animals with their respective standard errors).....	165
6.4	The apparent degradation rates of RNA, DNA and the absorption rate of PB (Y , % of the first segment) from difference intestinal segments(X , cm) of swamp buffalo and KK cattle.....	166
6.5	The quantity of fresh digesta (g) and PB (μmol) in each segment of small intestine of swamp buffaloes and KK cattle ...	169
7.1	Purine ($\mu\text{mol/g}$ DM) and N (mg/g DM) contents in mixed rumen microbial population of swamp buffaloes and KK cattle	177
7.2	Purine content and N contents in rumen microbial population of swamp buffaloes and KK cattle.....	179

LIST OF FIGURES

Figure	Page
2.1 Summarize of the purine metabolism in ruminants (Nolan, 1999)....	25
3.1 Chromatogram of standard PD sample obtained using HPLC, Peak 1= allantoin; 2=uric acid; 3=creatinine; 4= hypoxanthine; 5= xanthine; 6= oxypurinol (internal standard).....	59
3.2 Chromatogram of PD from urine sample obtained using HPLC, Peak 1= allantoin; 2= uric acid; 3= creatinine; 4= oxypurinol.....	59
3.3 Excretion of urinary purine derivatives in swamp buffaloes and KK cattle during last 5 days pre-fasting, and 6 days of a fasting period.....	62
3.4 Relationship between urinary PD excretion in Swamp buffaloes, KK cattle and digestible OM intake (DOMI).....	78
4.1 KK cattle fitted with a ruminal and duodenal cannula.....	87
4.2 Schedule of experimental procedure.....	89
4.3 The PB solution was continuously infused into duodenal cannulae via a vinyl tube.....	92
4.4 Chromatogram of standard PB sample obtained using HPLC, Peak 1= guanine, 2=allapurinol (internal standard) and 3=adenine.....	96
4.5 Chromatogram of PB from duodenal digesta sample obtained using HPLC, Peak 1= guanine, 2=allapurinol (internal standard) and 3=adenine.....	96
4.6 Urinary excretion of PD in relation to purine flows to duodenal in KK cattle given a duodenal infusion of PB.....	100
4.7 Four buffaloes were kept in individual metabolism crates.....	109
4.8 The PB solution was infused into the duodenal catheter via vinyl tube.....	111
4.9 Urinary excretion of PD in relation to purine flows to duodenum in swamp buffaloes given a duodenal infusion of PB.....	116

5.1	The concentration of plasma PD labeled with ^{14}C (% of the concentration of the injected) in swamp buffaloes.....	131
5.2	The concentration of plasma PD labeled with ^{14}C (% of the concentration of the injected) in KK cattle.....	132
5.3	The recovery of tracer $^{14}\text{C}_1$ (allantoin plus uric acid and other form) in 2 feeding levels of 2 buffaloes	136
5.4	The recovery of tracer $^{14}\text{C}_1$ (allantoin plus uric acid and other form) in 2 feeding levels of 2 KK cattle	136
5.5	The concentration of plasma PD labeled ^{14}C (% of injected) in 4 swamp buffaloes.....	144
5.6	The recovery of PD labeled ^{14}C in urine of swamp buffaloes.....	147
6.1	The small intestine of cattle and swamp buffalo tied with string at approximately 60 cm intervals.....	155
6.2	The RNA degradation in small intestine of buffaloes and KK cattle.....	161
6.3	The DNA degradation in small intestine of buffaloes and KK cattle.....	162
6.4	The purine bases absorbed in small intestine from buffaloes and KK cattle.....	167
7.1	Procedure to isolate solid associated microbial (SAM) and liquid associated microbes (LAM) from rumen contents, (modified procedure of IAEA- TECDOC, 1997).....	175

LIST OF ABBREVIATIONS

μCi	Micro Curi
μmol	Micromol
ADF	Acid Detergent Fiber
ADG	Average Daily Gain
AEP	2-aminoethylphosphonic acid
ATP	Adenine Triphosphate
BCP	Bacterial plus Protozoal Crude Protein
BW	Body Weight
C	Creatinine
°C	Degree Celsius
CF	Crude Fiber
CMC	Carboxy methyl Cellulose
Con	Concentrate
CP	Crude Protein
Cr-EDTA	Chromium-Ethylene diamine Tetra acetic acid
CV	Coefficient of Variation
DAPA	Diaminopimelic acid
DDMI	Digestible Dry Matter Intake
DM	Dry Matter
DMI	Dry Matter Intake
DNA	Deoxyribonucleic acid
DOM	Digestible Organic Matter
DOMI	Digestible Organic Matter Intake
DOMR	Digestible Organic Matter apparently fermented in the Rumen
DOMS	Digestible Organic Matter in the Stomach
dpm	Disintegrations per minute
EE	Ether Extract
FB	Fluid Associated Bacteria
FM	Fluid Associated Microbes
GE	Gross Energy
GFR	Glomerular Filtration Rate
h	Hour
HPLC	High Pressure Liquid Chromatography
IMP	5-inosine-monophosphate
Kg	Kilogram
Kg ^{0.75}	Metabolic Body Weight
KJ	Kilo-Joule
KK Cattle	Kedah Kelantan Cattle
L	Liter